

CONTINUED FRACTIONS, STATISTICS, AND GENERALIZED PATTERNS

TOUFIK MANSOUR

LABRI, Université Bordeaux I,
351 cours de la Libération, 33405 Talence Cedex
toufik@labri.fr

ABSTRACT

Recently, Babson and Steingrímsson (see [BS]) introduced generalized permutations patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation.

Following [BCS], let $e_k\pi$ (respectively; $f_k\pi$) be the number of the occurrences of the generalized pattern 12-3-...- k (respectively; 21-3-...- k) in π . In the present note, we study the distribution of the statistics $e_k\pi$ and $f_k\pi$ in a permutation avoiding the classical pattern 1-3-2.

Also we present an applications, which relates the Narayana numbers, Catalan numbers, and increasing subsequences, to permutations avoiding the classical pattern 1-3-2 according to a given statistics on $e_k\pi$, or on $f_k\pi$.

1. INTRODUCTION

Permutation patterns: let $\alpha \in S_n$ and $\tau \in S_k$ be two permutations. We say that α *contains* τ if there exists a subsequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $(\alpha_{i_1}, \dots, \alpha_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called a *pattern* or (*classical pattern*). We say α *avoids* τ , or is τ -*avoiding*, if such a subsequence does not exist. The set of all τ -avoiding permutations in S_n is denoted $S_n(\tau)$. For an arbitrary finite collection of patterns T , we say that α avoids T if α avoids any $\tau \in T$; the corresponding subset of S_n is denoted $S_n(T)$.

While the case of permutations avoiding a single pattern has attracted much attention, the case of multiple pattern avoidance remains less investigated. In particular, it is natural, as the next step, to consider permutations avoiding pairs of patterns τ_1, τ_2 . This problem was solved completely for $\tau_1, \tau_2 \in S_3$ (see [SS]), for $\tau_1 \in S_3$ and $\tau_2 \in S_4$ (see [W]), and for $\tau_1, \tau_2 \in S_4$ (see [B, K] and references therein). Several recent papers [CW, MV1, Kr, MV2, MV3] deal with the case $\tau_1 \in S_3, \tau_2 \in S_k$ for various pairs τ_1, τ_2 . Another natural question is to study permutations avoiding τ_1 and containing τ_2 exactly t times. Such a problem for certain $\tau_1, \tau_2 \in S_3$ and $t = 1$ was investigated in [R], and for certain $\tau_1 \in S_3, \tau_2 \in S_k$ in [RWZ, MV1, Kr, MV2]. The tools involved in these papers include continued fractions, Chebyshev polynomials, and Dyck paths.

Generalized permutation patterns: in [BS] Babson and Steingrímsson introduced generalized permutation patterns that allow the requirement that two

adjacent letters in a pattern must be adjacent in the permutation. This idea in introducing these patterns was study of Mahonian statistics.

We write a classical pattern with dashes between any two adjacent letters of the pattern, say 1324, as 1-3-2-4, and if we write, say 24-1-3, then we mean that if this pattern occurs in permutation π , then the letters in the permutation π that correspond to 2 and 4 are adjacent (see [C]). For example, the permutation $\pi = 35421$ has only two occurrences of the pattern 23-1, namely the subsequences 352 and 351, whereas π has four occurrences of the pattern 2-3-1, namely the subsequences 352, 351, 342 and 341.

Claesson [C] gave a complete answer for the number of permutations avoiding any single 3-letters generalized pattern with exactly one adjacent pair of letters. Later, Claesson and Mansour [CM] presented a complete solution for the number of permutations avoiding any double 3-letters generalized patterns with exactly one adjacent pair of letters. Besides, Kitaev [Ki] investigate simultaneous avoidance of two or more 3-letters generalized patterns without internal dashes.

On the other hand, Robertson, Wilf and Zeilberger [RWZ] showed a simple continued fraction that records the joint distribution of the patterns 1-2 and 1-2-3 on 1-3-2-avoiding permutations. Recently, generalization of this theorem given, by Mansour and Vainshtein [MV1], by Krattenthaler [Kr], by Jani and Rieper [JR], and by Brändén, Claesson and Steingrímsson [BCS].

Mansour [M, Th. 2.1 and Th. 2.9] presented an analog of this theorem ([MV1]) by replace generalized patterns with classical patterns.

In the present note, we generalize [M, Th. 2.1] and [M, Th. 2.9] and give an analog for [BCS] by replace generalized patterns with classical patterns. This by use the same argument proof of the main results in [BCS] with a simple changes. In the last section, we present an applications for our results.

2. MAIN RESULTS

For all $k \geq 1$, we denote by $a_k(\pi)$ the number of the occurrences of the pattern 1-2-3-...- k in π . In [BCS, Th. 1] proved the following.

Theorem 2.1. (P. Brändén, A. Claesson, and E. Steingrímsson, [BCS, Th. 1])

The $\sum_{\pi \in \mathcal{S}(1-3-2)} \prod_{k \geq 1} x_k^{a_k(\pi)}$ is given by the following continued fraction:

$$1 - \frac{1}{x_1^{(0)}} \quad 1 - \frac{x_1^{(1)} x_2^{(1)}}{x_1^{(0)} x_2^{(1)}} \quad 1 - \frac{x_1^{(2)} x_2^{(1)} x_3^{(2)}}{x_1^{(0)} x_2^{(1)} x_3^{(2)}} \quad \ddots$$

in which the $(n+1)$ st numerator is $\prod_{k=0}^n x_{k+1}^{(n)}$.

For all $k \geq 3$, we denote by $e_k(\pi)$ (respectively; $f_k(\pi)$) the number of the occurrences of the generalized pattern 12-3-...- k (respectively; 21-3-...- k) in π . Besides,

$e_2(\pi)$, $f_2(\pi)$, and $e_1(\pi) = f_1(\pi)$ are denote the number of occurrences of the pattern 12, 21 and 1, respectively, and extended that by $e_0(\pi) = f_0(\pi) = 1$.

Now let us give an analog for Theorem 2.1 by used generalized patterns with simply changes in the proof of Theorem 2.1.

Theorem 2.2. *The $\sum_{\pi \in \mathcal{S}(1-3-2)} \prod_{k \geq 1} x_k^{e_k(\pi)}$ is given by the following continued fraction:*

$$\frac{1}{1 - x_1 + x_1 x_2^{(0)} - \frac{x_1 x_2^{(0)}}{1 - x_1 + x_1 x_2^{(1)} x_3^{(1)} - \frac{x_1 x_2^{(1)} x_3^{(1)}}{1 - x_1 + x_1 x_2^{(2)} x_3^{(1)} x_4^{(2)} - \frac{x_1 x_2^{(2)} x_3^{(1)} x_4^{(2)}}{\ddots}}}}$$

in which the $(n+1)$ st numerator is $x_1 \cdot \prod_{k=0}^n x_{k+2}^{(k)}$.

Proof. Let π any nonempty permutation avoiding 1-2-3 such that $\pi = (\pi', n, \pi'')$ and $j = \pi^{-1}(n)$; so every element in π' is greater than every element in π'' . Thus $\pi', \pi'' \in \mathcal{S}(1-3-2)$, so

$$e_k(\pi) = \begin{cases} e_k(\pi') + e_{k-1}(\pi') + e_k(\pi''), & \text{for all } k \geq 3; \\ e_2(\pi') + e_2(\pi'') + \delta_{\pi', \emptyset}, & k = 2; \\ e_1(\pi') + e_1(\pi'') + 1, & k = 1 \end{cases},$$

where $\delta_{\pi', \emptyset}$ is the Kronecker delta.

It follows that the generating function

$$C(x_1, x_2, \dots) = \sum_{\pi \in \mathcal{S}(1-3-2)} \prod_{k \geq 1} x_k^{e_k(\pi)}$$

satisfies (see [M, Th. 2.1])

$$C(x_1, x_2, \dots) = 1 + x_1 C(x_1, x_2, \dots) + x_1 x_2 C(x_1, x_2, \dots) (C(x_1, x_2 x_3, x_3 x_4, \dots) - 1),$$

which means that ,

$$C(x_1, x_2, x_3, x_4, \dots) = \frac{1}{1 - x_1 + x_1 x_2 - x_1 x_2 C(x_1, x_2 x_3, x_3 x_4, \dots)}$$

and the theorem follows by induction. $\square$

Similarly, it is easy to see

$$f_k(\pi) = \begin{cases} f_k(\pi') + f_{k-1}(\pi') + f_k(\pi''), & \text{for all } k \geq 3; \\ f_2(\pi') + f_2(\pi'') + \delta_{\pi'', \emptyset}, & k = 2; \\ f_1(\pi') + f_1(\pi'') + 1, & k = 1 \end{cases},$$

where $\delta_{\pi'', \emptyset}$ is the Kronecker delta. So by using [M, Th. 2.9] and the argument proof of Theorem 2.1 we get the following.

Theorem 2.3. *The $\sum_{\pi \in \mathcal{S}(1-3-2)} \prod_{k \geq 1} x_k^{f_k(\pi)}$ is given by the following continued fraction:*

$$1 - \frac{x_1}{x_1 x_2^{(0)} - \frac{1}{1 - \frac{x_1}{x_1 x_2^{(1)} x_3^{(1)} - \frac{1}{1 - \frac{x_1}{x_1 x_2^{(2)} x_3^{(2)} x_4^{(2)} - \frac{1}{\ddots}}}}}}.$$

Following [BCS], we define $\mathcal{A}$ be the ring of all infinite matrices with finite number of non zero entries in each row, that is,

$$\mathcal{A} = \{A : \mathcal{N} \times \mathcal{N} \rightarrow \mathcal{Z} | \forall n (A(n, k) = 0 \text{ for almost every } k)\},$$

with multiplication defined by $(AB)(n, k) = \sum_{i \geq 1} A(n, i)B(i, k)$. With each $A \in \mathcal{A}$ we now associate a family of statistics $\{< \mathbf{q}, A_k >\}_{k \geq 1}$ defined on $\mathcal{S}(1-3-2)$, where $\mathbf{q} = (q_1, q_2, q_3, \dots)$ and

$$< \mathbf{q}, A_k > = \sum_{i \geq 1} A(i, k) q_i.$$

Following [BCS], let us define mathematical objects with respect A as follows. Let $\mathbf{q} = (q_1, q_2, \dots)$ where the q_i are indeterminates; for each $A \in \mathcal{A}$ and $\pi \in \mathcal{S}(1-3-2)$ we define three objects as follows:

The *weight* $\eta(\pi, A; \mathbf{q})$, the *weight* $\mu(\pi, A; \mathbf{q})$, and the *weight* $\nu(\pi, A; \mathbf{q})$ of π with respect A , by

$$\eta(\pi, A; \mathbf{q}) = \prod_{k \geq 1} q_k^{< \mathbf{a}, A_k > \pi}, \quad \mu(\pi, A; \mathbf{q}) = \prod_{k \geq 1} q_k^{< \mathbf{e}, A_k > \pi}, \quad \nu(\pi, A; \mathbf{q}) = \prod_{k \geq 1} q_k^{< \mathbf{f}, A_k > \pi},$$

respectively, where $\mathbf{a} = (a_1, a_2, \dots)$, $\mathbf{e} = (e_1, e_2, \dots)$ and $\mathbf{f} = (f_1, f_2, \dots)$.

The *generating function with respect* A of the, statistics $\{< \mathbf{a}, A_k >\}_{k \geq 1}$, statistics $\{< \mathbf{e}, A_k >\}_{k \geq 1}$, and statistics $\{< \mathbf{f}, A_k >\}_{k \geq 1}$ by

$$F_A(\mathbf{q}) = \sum_{\pi \in \mathcal{S}(1-3-2)} \eta(\pi, A; \mathbf{q}), \quad G_A(\mathbf{q}) = \sum_{\pi \in \mathcal{S}(1-3-2)} \mu(\pi, A; \mathbf{q}), \quad H_A(\mathbf{q}) = \sum_{\pi \in \mathcal{S}(1-3-2)} \nu(\pi, A; \mathbf{q}),$$

respectively.

The *continued fractions with respect* A , by

$$C_A(\mathbf{q}) = \frac{1}{\prod_{k \geq 0} q_{k+1}^{A(1,k)}},$$

$$1 - \frac{\prod_{k \geq 0} q_{k+1}^{A(2,k)}}{1 - \frac{\prod_{k \geq 0} q_{k+1}^{A(3,k)}}{\ddots}},$$

$$D_A(\mathbf{q}) = \frac{1}{1 - q_1 + q_1 \prod_{k \geq 1} q_{k+1}^{A(1,k)} - \frac{q_1 \prod_{k \geq 1} q_{k+1}^{A(1,k)}}{1 - q_1 + q_1 \prod_{k \geq 1} q_{k+1}^{A(2,k)} - \frac{q_1 \prod_{k \geq 1} q_{k+1}^{A(2,k)}}{\ddots}}},$$

and by

$$E_A(\mathbf{q}) = 1 - \frac{q_1}{q_1 \prod_{k \geq 1} q_{k+1}^{A(1,k)} - \frac{1}{1 - \frac{q_1}{q_1 \prod_{k \geq 1} q_{k+1}^{A(2,k)} - \frac{1}{\ddots}}}}.$$

As a remark (see [BCS, p. 3]): definition of the ring $\mathcal{A}$, and the fact that $a_i(\pi) = e_i(\pi) = f_i(\pi) = 0$ for all $i = |\pi| + 1, |\pi| + 2, \dots$ yields the product in the weight is finite.

The second step in [BCS], proved the following.

Theorem 2.4. (P. Brändén, A. Claesson, and E. Steingrímsson, [BCS, Th. 2])
For $A \in \mathcal{A}$, $F_A(\mathbf{q}) = C_{BA}(\mathbf{q})$ where $B = \left[\begin{smallmatrix} i \\ j \end{smallmatrix} \right]$, and conversely $C_A(\mathbf{q}) = F_{B^{-1}A}(\mathbf{q})$.

By above definitions with using the proof of Theorem 2.4 we get an analog of Theorem 2.4 for the statistics e_k and f_k as follows.

Theorem 2.5. Let $A \in \mathcal{A}$; then

$$\begin{aligned} G_A(\mathbf{q}) &= D_{BA}(\mathbf{q}), & H_A(\mathbf{q}) &= E_{BA}(\mathbf{q}), \\ D_A(\mathbf{q}) &= G_{B^{-1}A}(\mathbf{q}), & E_A(\mathbf{q}) &= H_{B^{-1}A}(\mathbf{q}), \end{aligned}$$

where $B = \left[\begin{smallmatrix} i \\ j \end{smallmatrix} \right]$, $\binom{n}{k} = 0$ for all $k > 0$ or $n > k$.

Proof. By using definitions we get that (here, we using the same argument proof in [BCS, Th. 2])

$$\begin{aligned} \mu(\pi, A; \mathbf{q}) &= \prod_{k \geq 1} q_k^{<\mathbf{e}, A_k>\pi} \\ &= \prod_{k \geq 1} \prod_{j \geq 1} q_k^{A(j,k)e_j(\pi)} \\ &= \prod_{j \geq 1} \left(\prod_{k \geq 1} q_k^{A(j,k)} \right)^{e_j(\pi)}. \end{aligned}$$

Let $x_{j+1} = \prod_{k \geq 1} q_k^{A(j,k)}$; Theorem 2.2 yields

$$\prod_{j \geq 1} x_{j+1}^{\binom{n-1}{j-1}} = \prod_{j \geq 1} \left(\prod_{k \geq 1} q_k^{A(j,k)} \right)^{\binom{n-1}{j-1}} = \prod_{k \geq 1} q_k^{((\binom{n-1}{0}), (\binom{n-1}{1}), (\binom{n-1}{2}), \dots), A_k}.$$

so, again, by definitions $G_A(\mathbf{q}) = D_{BA}(\mathbf{q})$. Observing that $B^{-1} = \left[(-1)^{i-j} \binom{i}{j} \right] \in \mathcal{A}$ we also obtain $D_A(\mathbf{q}) = G_{B^{-1}A}(\mathbf{q})$.

Similarly, we have $E_{BA}(\mathbf{q}) = H_A(\mathbf{q})$ and $E_A(\mathbf{q}) = H_{B^{-1}A}(\mathbf{q})$. $\square$

Remark 2.6. The general approach which described in [BCS] on the statistics $a_k(\pi)$ its work with others statistics. Its work on the statistics $e_k(\pi)$ and on the statistics $f_k(\pi)$. So the natural question to ask is the following: If there any descriptions for all the statistics such this approach will work? Here we failed to give an answer for this.

3. APPLICATION

In the current section, we present an examples for application of Theorem 2.5. Some of these examples are related known continued fraction to the statistics e_k or f_k , but others relate these statistics to others combinatorial objects.

3.1. Narayana numbers. Let $N(n, k) = \frac{1}{n} \binom{n}{k} \binom{n}{k+1}$ be the Narayana numbers, and the corresponding generating function for the Narayana numbers we denote by $N(x, t)$. Then

$$N(x, t) := \sum_{n, k \geq 0} N(n, k) x^k t^n = 1 + xtN^2(x, t) - xtN(x, t) + tN(x, t).$$

This allow us to express $N(x, t)$ as a continued fraction:

$$N(x, t) = \frac{1}{1 - \frac{t}{1 - \frac{tx}{1 - \frac{t}{1 - \frac{tx}{\ddots}}}}}$$

Proposition 3.1. *The number $N(n, k)$ equals the number of permutations $\pi \in S_n(132)$ with $e_2(\pi) = k$.*

Proof. Let $A(n, k) = \delta_{(n, k), (1, 1)} + \delta_{(n, k), (2, 2)}$ where δ is Kronecker delta, so by applying Theorem 2.5 we get that

$$N'(x, t) := \sum_{\pi \in S(1-3-2)} x^{e_2(\pi)} t^{|\pi|} = \frac{1}{1 - t - xt + xtN'(x, t)} = \frac{1}{1 - t - xt + \frac{xt}{1 - t - xt + \frac{xt}{\ddots}}},$$

so $N'(x, t)$ satisfies the same functional equation as $N(x, t)$, hence $N'(x, t) = N(x, t)$. $\square$

Again, the same argument work also to statistics on f_k as follows.

Proposition 3.2. *The number $N(n, k)$ equals the number of permutations $\pi \in S_n(132)$ with $f_2(\pi) = k$.*

Proof. Let $A(n, k) = \delta_{(n, k), (1, 1)} + \delta_{(n, k), (2, 2)}$ where δ is Kronecker delta, so by applying Theorem 2.5 we get that

$$N''(x, t) := \sum_{\pi \in S(1-3-2)} x^{f_2(\pi)} t^{|\pi|} = 1 - \frac{t}{xt - \frac{1}{1 - tN''(x, t)}} = 1 - \frac{t}{xt - \frac{1}{1 - \frac{t}{xt - \frac{1}{1 - \frac{t}{\ddots}}}}},$$

so $N''(x, t)$ satisfies the same functional equation as $N(x, t)$, hence $N''(x, t) = N(x, t)$. $\square$

3.2. Increasing subsequences. . As consequence of [BCS], we define as follows. The subsequence $\pi_{i_1} \pi_{i_1+1} \pi_{i_2} \dots \pi_{i_k}$ ($k \geq 2$) of π is called *2-increasing subsequence* if $\pi_{i_j} < \pi_{i_{j+1}}$, $i_j < i_{j+1}$ and $i_1 + 1 < i_2$. Hence, the total number of 2-increasing subsequences in a permutation is counting by $e_2 + e_3 + \dots$. An application of

Theorem 2.5 gives the following continued fraction for the distribution of $e_2 + e_3 + \dots$:

$$\sum_{\pi \in \mathcal{S}(1-3-2)} x^{e_2(\pi) + e_3(\pi) + \dots} t^{|\pi|} = \frac{1}{1 - t(1 - x) - \frac{xt}{1 - t(1 - x^2) - \frac{x^2t}{1 - t(1 - x^4) - \frac{x^4t}{\ddots}}}}.$$

The subsequence $\pi_{i_1} \pi_{i_1+1} \pi_{i_2} \dots \pi_{i_k}$ ($k \geq 2$) of π is called *almost 2-increasing subsequence* if $\pi_{i_j} < \pi_{i_{j+1}}$, $i_j < i_{j+1}$ for $j = 2, 3, \dots, k-1$, $i_1 + 1 < i_2$ and $\pi_{i_1+1} < \pi_{i_1}$. Hence, the total number of almost 2-increasing subsequences in a permutation is counting by $f_2 + f_3 + \dots$. An application of Theorem 2.5 gives the following continued fraction for the distribution of $f_2 + f_3 + \dots$:

$$\sum_{\pi \in \mathcal{S}(1-3-2)} x^{f_2(\pi) + f_3(\pi) + \dots} t^{|\pi|} = 1 - \frac{t}{xt - \frac{1}{1 - \frac{t}{x^2t - \frac{1}{1 - \frac{t}{x^4t - \frac{1}{1 - \frac{t}{x^8t - \ddots}}}}}}}}.$$

3.3. Catalan numbers. The n th Catalan number is given by $C_n = \frac{1}{n+1} \binom{2n}{n}$, and the corresponding generating function is given by $C(x) = \frac{1 - \sqrt{1-4x}}{2x}$. Theorem 2.5 yields for the statistic $s = 0$ ($s(\pi) = 0$ for all π) the following. The generating function $C(x)$ for the number of permutations avoiding 1-3-2 can be expressed, again, in terms of continued fractions:

$$C'(x) := \frac{1}{1 - \frac{x}{1 - \frac{x}{1 - \frac{x}{\ddots}}}}, \quad C''(x) := 1 - \frac{x}{x - \frac{1}{1 - \frac{x}{x - \frac{1}{\ddots}}}}.$$

In the above two cases $C'(x) = C''(x) = C(x)$.

REFERENCES

- [B] M. BÖNA, The permutation classes equinumerous to the smooth class, *Electron. J. Combin.* **5** (1998), #R31
- [BCS] P. BRÄNDÉN, A. CLAEISSON, AND E. STEINGRIMSSON, Continued fractions and increasing subsequences in permutations, to appear.
- [BS] E. BABSON AND E. STEINGRIMSSON, Generalized permutation patterns and a classification of the Mahonian statistics, *Séminaire Lotharingien de Combinatoire*, B44b:18pp, (2000).
- [C] A. CLAEISSON, Generalised pattern avoidance, *European Journal of Combinatorics*, **22** (2001) 961–973.
- [CM] A. CLAEISSON AND T. MANSOUR, Permutations avoiding a pair of generalized patterns of length three with exactly one dash, preprint CO/0107044.
- [CW] T. CHOW AND J. WEST, Forbidden subsequences and Chebyshev polynomials, *Discr. Math.*, **204** (1999) 119–128.
- [JR] M. JANI AND R.G. RIEPER, Continued fractions and catalan problems, *Elect. J. Combin.* **7** (2000) #R1.
- [Ki] S. KITAEV, Multi-Avoidance of generalized patterns, to appear.

- [Kn] D. KNUTH, The Art of Computer Programming, volume 3, Addison Wesley, Reading, MA 1973.
- [Kr] C. KRATTENTHALER, Permutations with restricted patterns and Dyck paths, (2000), preprint CO/0002200.
- [K] D. KREMER, Permutations with forbidden subsequences and a generalized Schröder number, *Disc. Math.* **218** (2000), 121–130.
- [LS] V. LAKSHMIBAI AND B. SANDHYA, Criterion for smoothness of Schubert varieties in $Sl(n)/B$, *Proc. Indian Acad. Sci.* **100** no. 1 (1990) 45–52.
- [LW] L. LORENTZEN AND H. WAADELAND, Continued fractions with applications, North-Holland 1992.
- [M] T. MANSOUR, Continued fractions and generalized patterns, preprint CO/0110037.
- [MV1] T. MANSOUR AND A. VAINSHTEIN, Restricted permutations, continued fractions, and Chebyshev polynomials, *Electron. J. Combin.*, **7** (2000) #R17.
- [MV2] T. MANSOUR AND A. VAINSHTEIN, Restricted 132-avoiding permutations, *Adv. Appl. Math.* **126** (2001), 258–269.
- [MV3] T. MANSOUR AND A. VAINSHTEIN, Layered restrictions and Chebyshev polynomials (2000), *Annals of Combinatorics*, to appear (2001), preprint CO/0008173.
- [N] J. NOONAN, The number of permutations containing exactly one increasing subsequence of length three, *Discr. Math.* **152** (1996) 307–313.
- [NZ] J. NOONAN AND D. ZEILBERGER, The enumeration of permutations with a prescribed number of “forbidden” patterns, *Adv. Appl. Math.* **17** (1996) 381–407.
- [SS] R. SIMION AND F. SCHMIDT, Restricted permutations *Europ. J. Comb.* **6** (1985) 383–406.
- [R] A. ROBERTSON, Permutations containing and avoiding 123 and 132 patterns, *Disc. Math. and Theo. Comp. Sci.* **3** (1999), 151–154.
- [RWZ] A. ROBERTSON, H. WILF, AND D. ZEILBERGER, Permutation patterns and continuous fractions, *Elec. J. Comb.* **6** no. 1 (1999) #R38.
- [W] J. WEST, Generating trees and the Catalan and Schröder numbers, *Discr. Math.* **146** (1995) 247–262.